

Written examination: 17. May 2020

Course name and number: **Introduction to Statistics (02323)**

Duration: 4 hours

Aids and facilities allowed: All

The questions were answered by

\_\_\_\_\_ (student number)

\_\_\_\_\_ (signature)

\_\_\_\_\_ (table number)

This exam consists of 30 questions of the “multiple choice” type, which are divided between 11 exercises. To answer the questions, you need to fill in the “multiple choice” form (6 separate pages) on CampusNet with the numbers of the answers that you believe to be correct.

5 points are given for a correct “multiple choice” answer, and  $-1$  point is given for a wrong answer. ONLY the following 5 answer options are valid: 1, 2, 3, 4, or 5. If a question is left blank or an invalid answer is entered, 0 points are given for the question. Furthermore, if more than one answer option is selected for a single question, which is in fact technically possible in the online system, 0 points are given for the question. The number of points needed to obtain a specific mark or to pass the exam is ultimately determined during censoring.

**The final answers should be given by filling in and submitting the form online via CampusNet. The table provided here is ONLY an emergency alternative. Remember to provide your student number if you do hand in on paper.**

<b>Exercise</b>	I.1	I.2	II.1	II.2	II.3	III.1	III.2	IV.1	IV.2	IV.3
<b>Question</b>	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
<b>Answer</b>										

<b>Exercise</b>	V.1	V.2	VI.1	VI.2	VI.3	VII.1	VII.2	VII.3	VIII.1	VIII.2
<b>Question</b>	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)
<b>Answer</b>										

<b>Exercise</b>	VIII.3	VIII.4	VIII.5	VIII.6	IX.1	IX.2	X.1	X.2	X.3	XI.1
<b>Question</b>	(21)	(22)	(23)	(24)	(25)	(26)	(27)	(28)	(29)	(30)
<b>Answer</b>										

The exam paper contains 23 pages.

Continue on page 2

**Multiple choice questions:** *Note that in each question, one and only one of the answer options is correct. Furthermore, not all the suggested answers are necessarily meaningful. Always remember to round your own result to the number of decimals given in the answer options before you choose your answer. Also remember that there may be slight discrepancies between the result of the book's formulas and corresponding built-in functions in R.*

### **Exercise I**

The characteristics of electrical components are not exactly as specified, e.g. if you buy a resistor then the resistance through it is not exactly as specified. In the production of electric circuits, it is of great interest not to get too much variation in the quality of the overall circuit. An example is the resistance through two parallel connected resistors, which is calculated by

$$R = \frac{1}{\frac{1}{R_1} + \frac{1}{R_2}}$$

where  $R_1$  is the resistance through one of the resistors and  $R_2$  through the other resistor. The resistance is measured in ohm.

Assume that  $R_1 \sim N(4, 0.2)$  and  $R_2 \sim N(2, 0.2)$ .

#### **Question I.1 (1)**

You buy 100  $R_1$  resistors - which can be assumed to be independent of each other. What is the probability that none of these has a resistance below 3 ohms?

- 1  1.27%
- 2  2.78%
- 3  13.9%
- 4  27.9%
- 5  42.4%

#### **Question I.2 (2)**

Calculate an estimate of the standard deviation of the total resistance  $R$  (the answer is rounded to two significant digits, tip: if you use simulation then remember to make sufficient repetitions to get a stable result)?

- 1  0.026
- 2  0.094

3  0.16

4  0.21

5  0.44

Continue on page 4


## Exercise II

In a computer system an optimization routine is used and the execution time for this routine is under investigation. The execution time is measured in hours and loaded into R with the following code:

```
x <- c(1.6, 2, 3.4, 4, 2.1, 0.6, 0.4, 0.4, 6, 0.4, 4.9, 2, 2, 4.6, 0.5,  
      3.4, 7.2, 10.5, 3.2, 1.3, 5.7, 1.9, 2.6, 2.5, 4.4, 1.8, 3.9, 6, 0.9)
```

### Question II.1 (3)

It is desirable to assess which distribution the outcomes in the sample could originate from. Therefore the following histogram has been generated of the observations in  $x$ :



On the basis of the information given, evaluate which of the following distributions is most likely to have generated the observations in the sample?

- 1  A normal distribution
- 2  A Poisson distribution
- 3  An exponential distribution
- 4  A  $t$ -distribution
- 5  A binomial-distribution

### Question II.2 (4)

Based on the sample, what is the estimate of the mean and standard deviation of the computation times?

- 1   $\hat{\mu} = 2.53$  and  $\hat{\sigma} = 1.66$
- 2   $\hat{\mu} = 3.36$  and  $\hat{\sigma} = 0.48$
- 3   $\hat{\mu} = 3.11$  and  $\hat{\sigma} = 2.37$
- 4   $\hat{\mu} = 1.98$  and  $\hat{\sigma} = 5.63$
- 5   $\hat{\mu} = 3.96$  and  $\hat{\sigma} = 2.81$

### Question II.3 (5)

You want to give a guarantee that the execution time is below a certain level, and therefore a confidence interval of the 90% quantile should be calculated. A function is defined in R to calculate it by:

```
q90 <- function(x){ quantile(x, prob=0.9, type=2) }
```

Which of the following R codes calculates a 95% percent non-parametric bootstrap confidence interval for the 90% quantile of the distribution of computation times?

- 1 

```
simsamples <- replicate(10000, sample(x, replace = TRUE))  
simmeans <- apply(simsamples, 2, q90)  
quantile(simmeans, c(0.05, 0.95))
```
- 2 

```
simsamples <- replicate(10000, sample(x, replace = FALSE))  
simmeans <- apply(simsamples, 2, q90)  
quantile(simmeans, c(0.025, 0.975))
```
- 3 

```
simsamples <- replicate(10000, sample(x, replace = FALSE))  
simmeans <- apply(simsamples, 2, q90)  
quantile(simmeans, c(0.05, 0.95))
```
- 4 

```
simsamples <- replicate(10000, sample(x, replace = TRUE))  
simmeans <- apply(simsamples, 2, q90)  
quantile(simmeans, c(0.1, 0.90))
```
- 5 

```
simsamples <- replicate(10000, sample(x, replace = TRUE))  
simmeans <- apply(simsamples, 2, q90)  
quantile(simmeans, c(0.025, 0.975))
```

Continue on page 6

### Exercise III

An NGO has 15 callers employed to recruit new members. Let  $X$  represent the number of members a single caller recruits during one working day. The number of new members each caller recruits in a day can be assumed to be independent of each other. From experience it is known that a good model for  $X$  is a binomial distribution, where the probability of getting a new member in a call is 7%. It is assumed that each caller can do 120 calls in one day.

#### Question III.1 (6)

What is the probability that a caller on a single day recruits more than 5 new members?

- 1  0.12
- 2  0.85
- 3  0.45
- 4  0.17
- 5  0.96

#### Question III.2 (7)

If  $Y$  is the total number of new members the 15 callers can recruit in a day, what is the mean and variance of  $Y$ ?

- 1   $E(Y) = 126$  og  $V(Y) = 10.8$
- 2   $E(Y) = 126$  og  $V(Y) = 41.9$
- 3   $E(Y) = 126$  og  $V(Y) = 43.5$
- 4   $E(Y) = 126$  og  $V(Y) = 102.4$
- 5   $E(Y) = 126$  og  $V(Y) = 117.2$

Continue on page 7

## Exercise IV

In Denmark, people often discuss whether it has been a good or a bad summer. The table below shows the average temperatures for the months May to September in the years 2014-2018:

	2014	2015	2016	2017	2018	Average
May	11.7	9.7	12.9	12.0	15.0	12.26
June	14.9	12.7	16.0	14.7	16.5	14.96
July	19.5	15.5	16.4	15.5	19.2	17.22
August	16.0	17.4	16.1	16.0	17.5	16.60
September	14.6	13.2	16.2	13.3	14.1	14.28
Average	15.34	13.70	15.52	14.30	16.46	15.01

To investigate if there is a difference between the years, the following R code has been run, where `year` is indicator for the years 2014-2018 and `temp` is the average temperature:

```
anova(fit <- lm(temp ~ year))  
  
## Analysis of Variance Table  
##  
## Response: temp  
##           Df Sum Sq Mean Sq F value Pr(>F)  
## year       4  23.4    5.85    1.15  0.36  
## Residuals 20 101.7    5.08
```

### Question IV.1 (8)

At the significance level  $\alpha = 0.05$  what is the conclusion from the R code above (by difference is meant significant difference in mean. Remember all parts of the answer must be correct)?

- 1  There is no difference in temperature between the years, since  $5.85 > 5.08$ .
- 2  There is a difference in temperature between the years, since  $0.36 > 0.05$ .
- 3  There is a difference in temperature between the years, since  $23.4 < 101.7$ .
- 4  There is no difference in temperature between the years, since  $0.36 > 0.05$ .
- 5  There is a difference in temperature between the years, since  $0.36 < 5$ .

### Question IV.2 (9)

The model used in the test above can be written as

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad \epsilon_{ij} \sim N(0, \sigma^2) \text{ and i.i.d.}$$

What is the estimate of  $\sigma^2$ ?

1   $\hat{\sigma}^2 = (23.4 + 101.7)/(4 + 20)$

2   $\hat{\sigma}^2 = 101.7/20$

3   $\hat{\sigma}^2 = \sqrt{101.7}$

4   $\hat{\sigma}^2 = (5.85 + 5.08)/(4 + 20)$

5   $\hat{\sigma}^2 = \sqrt{5.08}$

**Question IV.3 (10)**

What is

$$\sum_{\text{all } j,i} (y_{ij} - \bar{y})^2$$

where  $y_{ij}$  represents the observed temperatures and  $\bar{y}$  is the average of all the temperatures?

1  106.78

2  10.93

3  29.25

4  1.15

5  125.1

Continue on page 9



**Exercise V**

A sample of  $n = 50$  observations was taken from a population. The population distribution is not known, but it is known that the mean  $\mu$  is equal to 0. Let the  $i$ 'th observation be represented by  $X_i$  and the sample mean by  $\bar{X}$ . The sample standard deviation is denoted by  $s$ .

**Question V.1 (11)**

You now want to get an idea of the distribution of the population, which of the following actions would be the most obvious to carry out?

- 1  Make a density histogram of the sample.
- 2  Make a scatter plot of the population.
- 3  Make a qq-normal plot of the sample.
- 4  Make a pie chart of the population.
- 5  Make a  $t$ -test.

**Question V.2 (12)**

The following transformation has been defined

$$Y = \frac{\bar{X} - \mu}{\frac{s}{\sqrt{n}}}$$

Which of the following distributions approximates the distribution of  $Y$ ?

- 1  An exponential distribution.
- 2  A  $t$ -distribution with 2 degrees of freedom.
- 3  A  $\chi^2$ -distribution.
- 4  An  $F$ -distribution.
- 5  A standard normal distribution.

Continue on page 10

### Exercise VI

In a research project on energy consumption in schools, it was investigated how students and teachers set the radiator thermostats in classrooms. Thermostat settings were recorded during a period with cold weather at a number of schools in randomly selected classrooms. It was predetermined that thermostats are well set if the setting is between 2 and 3 on all radiators in a room, as it otherwise indicates under or oversized radiators. Besides, it's not desirable for thermostats to be set differently in a room, as this results in inferior comfort and poorer return water cooling.

The following observations were made during the period:

	School 1	School 2	School 3	School 4
Not-well set	18	11	22	9
Well set	38	36	15	12

Hence, at School 2 there were 11 out of 47 rooms in which the thermostats were not well set.

#### Question VI.1 (13)

What is the 95% confidence interval for the proportion of thermostats that were not set well at School 1 (note that the result from the R functions and the books formula may be slightly different, but both results are always closest to the correct answer)?

1  [0.07, 0.17]

2  [0.20, 0.44]

3  [0.26, 0.53]

4  [0.05, 0.22]

5  [0.18, 0.51]

#### Question VI.2 (14)

It was planned to compare schools to investigate if there were differences in practice of thermostat setting at the schools. Under the null hypothesis of no difference, then what is the expected number of not-well set thermostats at School 3?

1   $e_{13} = 37 \cdot \frac{60}{161} = 13.8$

2   $e_{13} = 15 \cdot \frac{22}{37} = 8.9$

3   $e_{13} = 60 \cdot \frac{124}{161} = 46.2$

4   $e_{13} = 22 \cdot \frac{22}{37} = 13.1$

5   $e_{13} = 15 \cdot \frac{124}{161} = 11.6$

**Question VI.3 (15)**

You want to investigate whether there was a difference in the proportion of not-well set thermostats at the four schools. Use a significance level of 1%. What will be the conclusion (both conclusion and argument must be correct)?

- 1  A difference between the schools cannot be detected, since the relevant test statistic is below the critical level of 15.4.
- 2  A difference between the schools can be detected, since the relevant test statistic is below the critical level of 11.3.
- 3  A difference between the schools cannot be detected, since the relevant test statistic is above the critical level of 0.0057.
- 4  A difference between the schools cannot be detected, since the relevant test statistic is below the critical level of 0.0057.
- 5  A difference between the schools can be detected, since the relevant test statistic is above the critical level of 11.3.

Continue on page 12

## Exercise VII

The following 3 questions deal with various statistical problems that may arise when treating water.

### Question VII.1 (16)

When controlling drinking water the quality is measured by regular water analysis. There is, of course, legislation on quality among other things requirements for concentrations of different substances. One requirement is that the conductivity of the water at the consumer's tap must not be more than  $2500 \mu\text{S}/\text{cm}$  at  $20^\circ\text{C}$ . If the conductivity is above this level, the concentration of salts is too high and the water is referred to as aggressive.

The conductivity of water at randomly selected consumers' taps has been measured. Let a measurement be represented by the stochastic variable  $X_i$ , which can be assumed to be normally distributed. Twenty independent measurements have been collected to determine the conductivity and the observed values are stored in the vector  $\mathbf{x}$  in R. You now want to test if the water on average is aggressive, and the following null hypothesis about the mean value,  $\mu$ , of the conductivity in the drinking water at the consumer's tap, is formulated

$$H_0 : \mu = 2500$$

with the alternative hypothesis

$$H_1 : \mu \neq 2500$$

The following from R is given:

```
t.test(x)

##
## One Sample t-test
##
## data: x
## t = 4.8527, df = 19, p-value = 0.0001106
## alternative hypothesis: true mean is not equal to 0
## 95 percent confidence interval:
## 704.1022 1772.1090
## sample estimates:
## mean of x
## 1238.106
```

Based on the result above and with a 5% significance level, which of the following statements is correct (both conclusion and reasoning must be correct)?

- 1  We reject  $H_0$  as the relevant confidence interval does not contain 0.

- 2  We accept  $H_0$ , since the sample average is within the relevant confidence interval.
- 3  We accept  $H_0$ , since the test statistic is greater than 1.96.
- 4  We reject  $H_0$ , since the  $p$ -value is 0.0001106.
- 5  We reject  $H_0$ , since 2500 is not within the relevant confidence interval.

### Question VII.2 (17)

At waterworks the water is purified by one of two methods, A or B, and the remaining concentration of a substance is measured. Measurements of remaining concentration are given in the stochastic variables  $Y_{A,i}$  and  $Y_{B,i}$  for methods A and B, respectively. It is of interest to investigate which of the two methods best purifies the water.  $Y_{A,i}$  and  $Y_{B,i}$  can be assumed to be normally distributed and their variances,  $\sigma_A^2$  and  $\sigma_B^2$ , can be assumed to be equal. For each of the two methods a sample of 20 observations is taken. We want to test the null hypothesis

$$H_0 : \mu_A = \mu_B$$

where the alternative hypothesis is

$$H_1 : \mu_A \neq \mu_B$$

Which of the following procedures is a correct approach?

- 1  A paired  $t$ -test with 19 degrees of freedom.
- 2  A paired  $t$ -test with 18 degrees of freedom.
- 3  A two-sample  $t$ -test with 38 degrees of freedom.
- 4  A two-sample  $t$ -test with 39 degrees of freedom.
- 5  A  $F$ -test testing variance homogeneity.

### Question VII.3 (18)

A new study is planned on the concentration of a substance in a drinking water drilling. We want to achieve a power of 90% to detect a mean value difference of 2 units from a given value. From experience it is known that the standard deviation is 3.5 units and you want to perform the test at a 5% significance level (remember that results from R functions may differ slightly from the result obtained with the book's formulas). How many observations must be taken to fulfil these requirements?

- 1  At least 15 observations.

- 2  At least 35 observations.
- 3  At least 48 observations.
- 4  At least 67 observations.
- 5  At least 102 observations.

Continue on page 15

### Exercise VIII

On 30 randomly selected summer days, corresponding values of the temperature at noon,  $x$  measured in degrees Celsius, and the number of ice creams sold in an ice cream chain,  $Y$ , have been recorded. The following model has been fitted in R:

$$Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i \quad \text{where } \varepsilon_i \sim N(0, \sigma^2) \text{ and i.i.d.}$$

The result of this is shown below:

```
summary(fit1)

##
## Call:
## lm(formula = y ~ x)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -161.24  -81.60  -46.14   103.83   249.85
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -2382.001    116.620  -20.43  <2e-16 ***
## x             230.703     5.083   45.38  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 124.7 on 28 degrees of freedom
## Multiple R-squared:  0.9866, Adjusted R-squared:  0.9861
## F-statistic: 2060 on 1 and 28 DF, p-value: < 2.2e-16
```

#### Question VIII.1 (19)

Based on the above R output, what is the estimate of the variance of the errors  $\hat{\sigma}^2$ ?

- 1  116.6
- 2   $116.6^2$
- 3  124.7
- 4   $124.7^2$
- 5   $(230.7/28)^2$

**Question VIII.2 (20)**

Based on the above R output, what is the prediction of the mean value of ice creams sold,  $\hat{y}_{\text{new}}$ , at  $x_{\text{new}} = 25^\circ \text{C}$ ?

- 1  231 ice creams.
- 2  2382 ice creams.
- 3  3386 ice creams.
- 4  5768 ice creams.
- 5  11535 ice creams.

**Question VIII.3 (21)**

Based on the above R output, what is the critical value for the test

$$H_0 : \beta_1 = 0$$

using a 1% significance level?

- 1  2.76
- 2  2.05
- 3  1.96
- 4  1.70
- 5  2.56

**Question VIII.4 (22)**

Which of the following statements about a prediction interval for  $Y_{\text{new}} = \hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}} + \varepsilon_{\text{new}}$  is not correct?

- 1  The prediction interval is wider than a corresponding confidence interval.
- 2  The width of the prediction interval depends on the sample size.
- 3  The prediction interval is symmetrical around the predicted value.
- 4  The width of the prediction interval depends on the value of  $x_{\text{new}}$ .



- 5  If the sample size becomes large enough, then the width of the prediction interval becomes 0.

### Question VIII.5 (23)

It is suspected that the linear model is not a correct model, so now you instead fit the model  $Y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \varepsilon_i$ . The model is fitted by (note that `x2` is `x` squared):

```
x2 <- x^2
fit2 <- lm(y ~ x + x2)
summary(fit2)

##
## Call:
## lm(formula = y ~ x + x2)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -193.67  -59.32  -25.73   64.96  263.87
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -159.3055   470.1439  -0.339    0.737
## x             24.9850    42.9277   0.582    0.565
## x2             4.5715     0.9502   4.811 5.06e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 93.15 on 27 degrees of freedom
## Multiple R-squared:  0.9928, Adjusted R-squared:  0.9922
## F-statistic: 1856 on 2 and 27 DF, p-value: < 2.2e-16
```

Based on this R output and with a 1% significance level, what can now be concluded about the relationship between ice cream sales and temperature (both conclusion and argument must both be correct)?

- 1  The relationship differs statistically significantly from a linear relationship, since  $\hat{\beta}_2$  is positive.
- 2  The relationship does not differ statistically significantly from a linear relationship, since the  $p$ -value for  $\hat{\beta}_1$  is above 1%.
- 3  The relationship differs statistically significantly from a linear relationship, since the  $p$ -value for  $\hat{\beta}_2$  is below 1%.


- 4  We cannot reject that the relationship is linear, since  $\hat{\beta}_2 < \hat{\beta}_1$ .
- 5  We cannot reject that the relationship is linear, since  $R^2 \approx 1$ .

**Question VIII.6 (24)**

In this exercise the result from the fit of the model from the previous question is investigated.

Below are shown two plots (A and B), where:

- $z$  is a new variable, which represent measurements of sunshine on a day (the unit doesn't matter).
- $x$  is the temperature on a day.
- `residuals(fit2)` is the residuals from the fit.



Which of the following statements is correct?

- 1  Figure B is used to examine whether the assumption of variance homogeneity is met.
- 2  Figure A is used to investigate whether the normal distribution assumption is met.
- 3  Figure B indicates a strong relationship between  $z$  og  $x$ .
- 4  Figure A can be used to investigate whether  $z$  should be included in the model.
- 5  Figure B can be used to check whether the relationship between  $x$  and  $Y$  is modelled correctly.

Continue on page 19

**Exercise IX**

The number of shooting stars per hour,  $X$ , is given by  $X \sim Po(3)$ , i.e. Poisson distributed with mean 3 shooting stars per hour.

**Question IX.1 (25)**

If one counts shooting stars for four hours, how many shooting stars might one expect to see (remember that the expectation value is equal to the mean)?

1  8

2  9

3  12

4  16

5  24

**Question IX.2 (26)**

Suppose one has just observed a shooting star. What is then the probability of waiting more than 10 minutes for the next shooting star?

1   $P(X = 0), X \sim Po(3)$

2   $P(X > 0), X \sim Po(3)$

3   $P(Y > 10), Y \sim Exp(3)$

4   $P(Y > \frac{10}{60}), Y \sim Exp(3)$

5   $P(Y > \frac{10}{6}), Y \sim Exp(3)$

Continue on page 20

## Exercise X

In a production of steel pipes one is interested in the diameter of the pipes. Therefore, a sample of 30 tubes is taken and the sample variance is calculated to be  $s^2 = 531 \text{ mm}^2$ .

### Question X.1 (27)

What is the 99% confidence interval for the standard deviation of the diameter of the tubes?

- 1  [18.4, 31.0]
- 2  [18.1, 30.6]
- 3  [310, 1079]
- 4  [17.2, 34.3]
- 5  [294, 1175]

### Question X.2 (28)

The diameter of the steel pipes is usually measured by one of two different measurement methods. It is now suspected that the two methods don't measure identically. 11 pipes are therefore randomly selected. Each pipe is now measured by both methods. The observation made with measurement Method 1 on pipe  $i$  is at position  $i$  in the vector  $\mathbf{x}$  and corresponding measurement with Method 2 at position  $i$  in the vector  $\mathbf{y}$ . The measurements by both methods can be assumed to be normally distributed.

The following analyses are now carried out in R:

```
t.test(x-y)

##
## One Sample t-test
##
## data:  x - y
## t = -2.541, df = 10, p-value = 0.0293
## alternative hypothesis: true mean is not equal to 0
## 95 percent confidence interval:
##  -2.409246 -0.158027
## sample estimates:
## mean of x
##  -1.28364
```

```
t.test(x,y)

##
## Welch Two Sample t-test
##
## data: x and y
## t = -0.1353, df = 20, p-value = 0.894
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
## -21.0683 18.5010
## sample estimates:
## mean of x mean of y
## 96.9018 98.1855
```

You want to test the null hypothesis that the two measurement methods have the same mean

$$H_0 : \mu_X = \mu_Y$$

against the alternative hypothesis that they are different

$$H_1 : \mu_X \neq \mu_Y$$

At a 5% significance level, which of the following possibilities is correct (both conclusion and argument must be correct)?

- 1  We reject  $H_0$ , since  $p < 0.05$ .
- 2  We cannot reject  $H_0$ , since  $p = 0.89$ .
- 3  We reject  $H_0$ , since the alternative hypothesis is different from 0.
- 4  We cannot reject  $H_0$ , since the test statistic,  $t_{obs}$ , is -0.14.
- 5  We reject  $H_0$ , since the difference between the sample averages is greater than 1.

### Question X.3 (29)

The company that produces the steel pipes is getting a new and faster machine for producing steel pipes. Regardless of the answer in the previous question, only measurement Method 1 is used below. The desired diameter of the steel pipes is 100 mm and the machine will be properly calibrated. You will plan a new experiment where you want to test the null hypothesis

$$H_0 : \mu_X = 100$$

against the alternative hypothesis

$$H_1 : \mu_X \neq 100$$

We use  $\hat{\sigma}_x^2 = 502.23$ , a significance level of 5% and the following R-code has been executed, since a sample size of  $n = 40$  is wanted:

```
power.t.test(n=40, sd=sqrt(502.23), power=0.9, type="one.sample")

##
##   One-sample t test power calculation
##
##           n = 40
##          delta = 11.8
##           sd = 22.4
##   sig.level = 0.05
##          power = 0.9
## alternative = two.sided
```

Based on the R code above, what can now be concluded before the experiment is performed (both argument and conclusion must be correct)?

- 1  If the true mean is 88.2 or lower, we have more than 90% chance of rejecting  $H_1$ .
- 2  If the true mean is 88.2 or lower, we have more than 90% chance of rejecting  $H_0$ .
- 3  If the true mean is 88.2 or lower, we will reject  $H_0$ .
- 4  If the true mean differs with less than 11.8 we accept  $H_0$ .
- 5  If the true mean differs with less than 11.8 we reject  $H_0$ .

Continue on page 23

**Exercise XI**

In the production of a particular type of plate, each plate has a 20% probability of having a flaw. A random sample of 10 plates is now taken.

**Question XI.1 (30)**

What is the probability that at most 3 plates in the sample have a flaw?

1  0.95

2  0.32

3  0.68

4  0.88

5  0.60

The exam is finished. Have a great summer!